arXiv:cond-mat/9807002v1 [cond-mat.stat-mech] 1 Jul 1998

Acceleration of quasi-particle modes in Bose-Einstein condensates

Karl-Peter Marzlin and Weiping Zhang
School of Mathematics, Physics, Computing and Electronics, Macquarie University, Sydney, NSW 2109, Australia

We analytically examine the dynamics of quasi-particle modes occuring in a Bose-Einstein condensate which is subject to a weak acceleration. It is shown that the momentum of a quasi-particle mode is squeezed rather than accelerated.

The achievement of Bose-Einstein condensation in magnetic traps [1] has led to a great interest in the physical properties of inhomogeneous atomic quantum gases. In the current research, one of the central topics is to study collective excitations of quasi-particle modes in Bose-Einstein condensates (BEC). The frequency spectrum of elementary excitations in a trapped ^{87}Rb BEC has been experimentally measured at JILA [2]. The propagation of sound waves in an excited BEC has been demonstrated at MIT [3]. Theoretical studies have revealed the characteristics of low-energy quasi-particle modes in such nonlinear systems [4].

In this paper we are interested in another aspect of collective excitations of BECs. We will study the dynamics of quasi-particle modes and show theoretically how they are manipulated by a weak homogeneous force. Since the dispersion relation of quasi-particle modes is much different from that of ordinary Schrödinger particles their dynamics will show up unusual effects like an acceleration-induced squeezing of position or momentum. A squeezing of the position may correspond to a localization of sound waves in quantum gases.

To initiate our idea, we start with a simplified model where a BEC trapped in a box of size L. In practice the box may be realized by a sufficiently anharmonic trap or an atomic cavity. We then add to this trapping potential a second, linear potential which describes the external homogeneous force. As a result, the complete potential becomes asymmetric and should incorporate an area where the potential is almost linear (see Fig. 1). A harmonic trap is not suited for this purpose since the homogeneous force then only leads to an overall displacement of the trap without changing its shape.

The time evolution of a BEC can be appropriately described by the time-dependent Gross-Pitaevskii equation,

$$i\hbar \frac{d\psi}{dt} = \left\{ \frac{p^2}{2M} + V(x) \right\} \psi + \kappa |\psi|^2 \psi \;, \tag{1}$$

where $\psi(x)$ is the normalized collective wavefunction of the BEC, M the atomic mass, and $\kappa := 4\pi\hbar^2 N a_{\rm scat}/M$. N is the number of condensed atoms and $a_{\rm scat} > 0$ the scattering length of the interaction between the atoms. $V(x)$ is an external potential.

The ground state $\psi_g(x,t) = \exp[-i\mu t/\hbar]\psi_g(x)$ of the trapped BEC is a stationary solution of Eq. (1), where μ denotes the chemical potential. $\psi_g(x)$ can be assumed to be real. For current experiments the nonlinear interaction between the atoms is relatively strong compared to the BEC's kinetic energy. It is then possible to neglect the kinetic term in Eq. (1) and to describe the ground state in the Thomas-Fermi approximation (TFA) [5],

$$\kappa |\psi_g|^2(x) = \mu - V(x) \;. \tag{2}$$

Of course this approximation is only valid in areas where the r.h.s. (right-hand-side) does not become small or negative.

Elementary excitations around the ground state ψ_g of Eq. (1) are described by writing the collective wavefunction as $\psi(x,t) = \exp[-i\mu t/\hbar]\{\psi_g(x) + \Theta(x,t)\}$ and expanding Eq. (1) up to terms linear in Θ. The resulting equation,

$$i\hbar \frac{d\Theta}{dt} = R\Theta + S\Theta^* \;, \tag{3}$$

with $R := p^2/(2M) + V - \mu + 2\kappa|\psi_g(x)|^2$ and $S := \kappa\psi_g^2$, has for time-independent R and S the formal solution

$$\begin{aligned} \Theta(x,t) = & \left\{ \cos[\hat\omega_+ t] - i(R+S)\frac{\sin[\hat\omega_+ t]}{\hat\omega_+} \right\} {\rm Re}\Theta_0 + \\ & i \left\{ \cos[\hat\omega_- t] - i(R-S)\frac{\sin[\hat\omega_- t]}{\hat\omega_-} \right\} {\rm Im}\Theta_0 \end{aligned} \tag{4}$$

($\Theta_0 := \Theta(x,0)$). The formal frequency operators

$$\hbar^2 \hat\omega_\pm := \sqrt{(R \mp S)(R \pm S)} \tag{5}$$

correspond to the spectrum elementary excitations which is usually calculated by using the Bogoliubov-DeGennes equations (see, e.g., Ref. [6]). We remark that one can rewrite Eq. (4) using $\hat\omega_+$ or $\hat\omega_-$ alone by exploiting relations like $(R-S)\cos[\hat\omega_- t] = \cos[\hat\omega_+ t](R-S)$.

In areas where the TFA is valid the operators $\hat\omega_\pm$ can be simplified to

$$\hbar\hat\omega_+ := \sqrt{\frac{\hat p^2}{2M}\left(\frac{\hat p^2}{2M} + 2\mu - 2V(\hat x) \right)} \;, \tag{6}$$

and a similar expression for $\hat\omega_-$. If we restrict our considerations to quasi-particle modes with small momentum we can neglect the second kinetic term in this equation and arrive at

$$\hbar\hat{\omega}_{+} := \sqrt{\hat{p}^{2}\left(c^{2} - V(\hat{x})/M\right)}\,, \tag{7}$$

where we have defined the velocity of sound $c := \sqrt{\mu/M}$. Obviously this approximation corresponds to the restriction to the linear part of the dispersion relation, $\hbar\hat{\omega}_{+} = c|\hat{p}|$, for elementary excitations in a homogeneous, free BEC ($V(x) = 0$).

Though Eq. (7) looks simple the square root on the r.h.s. is operator-valued and in general difficult to analyze. We therefore restrict our analysis to one spatial dimension and consider an accelerating potential $V(x) = -Max$ with $aL \ll c^{2}$, where a is the acceleration and L the length of the condensate. We then can expand $\hat{\omega}_{+}$ in a, leading to

$$\begin{aligned}
\hbar\hat{\omega}_{+} &= |\hat{p}|\left(1 + \frac{a\hat{x}}{2c^{2}} - \frac{a^{2}\hat{x}^{2}}{8c^{4}}\right) - i\frac{\hbar a}{4c^{2}}\mathrm{sgn}(\hat{p}) \\
&\qquad + i\frac{\hbar a^{2}}{8c^{4}}\mathrm{sgn}(\hat{p})\hat{x} + \frac{3\hbar^{2}a^{2}}{32c^{4}}|\hat{p}|^{-1} + O(a^{3}) \\
\hbar\hat{\omega}_{-} &= |\hat{p}|\left(1 + \frac{a\hat{x}}{2c^{2}} - \frac{a^{2}\hat{x}^{2}}{8c^{4}}\right) + i\frac{3\hbar a}{4c^{2}}\mathrm{sgn}(\hat{p}) \\
&\qquad - i\frac{3\hbar a^{2}}{8c^{4}}\mathrm{sgn}(\hat{p})\hat{x} + \frac{3\hbar^{2}a^{2}}{32c^{4}}|\hat{p}|^{-1} + O(a^{3})
\end{aligned} \tag{8}$$

In the derivation we have used the commutation relation $[\hat{x},\ \mathrm{sgn}(\hat{p})] = 2i\hbar\mathcal{P}_{0}$, where $\mathcal{P}_{0}$ is the projection operator on the state $p = 0$.

Some caution is needed when expanding the square root of an operator as in Eq. (8). Already Eq. (7) is only meaningful if $c^{2} - V/M = c^{2} + ax$ is larger than zero, in agreement with the assumptions of the TFA. Hence our approach is only useful for wavepackets that are sufficiently localized inside the interval $x \in [-c^{2}/a, -c^{2}/a + L]$. In addition we have to demand that the quasi-particle momentum is not too small so that its De-Broglie-wavelength is not larger than L. This becomes evident in the expansion (8) which diverges for $p \to 0$. Mathematically the inverse of an operator with eigenvalue zero cannot be defined. However, the inverse makes sense on a subspace that excludes this eigenvalue, and our restriction to sufficiently large p ensures that we are working in this subspace. Our approach therefore allows us to study wavepackets which are localized in phase space. For simplicity we will assume that the wavepacket is localized around some momentum $p_{0} > 0$ in momentum space so that we can set the operator $\mathrm{sgn}(\hat{p})$ equal to one. This causes no loss of generality as any wavepacket subject to the constraints given above can be linearly split into a part with $p > 0$ and one with $p < 0$. The latter case can be treated analogously to the case $p > 0$ which we will consider now.

Having derived an explicit expression for the frequency operators we can return to Eq. (4) and consider the evolution of a localized elementary excitation. Obviously the time dependence of Θ is essentially determined by the four exponentials $\exp[\pm i\hat{\omega}_{\pm}t]$. Since $\hat{\omega}_{-}$ is very similar to $\hat{\omega}_{+}$ we will only study the exponentials of $\hat{\omega}_{+}$ which can be decomposed as

$$\begin{aligned}
e^{-i\hat{\omega}_{+}t} &= \exp\left\{-\frac{at}{4c} + \frac{a^{2}t^{2}}{16c^{2}}\right\}\exp\left\{-i\frac{ct}{\hbar}\left(1 + \frac{at}{4c}\right)\hat{p}\right\} \\
&\quad \times \exp\left\{-i\frac{at}{2c\hbar}\left(1 - \frac{at}{4c}\right)\hat{p}\hat{x}\right\}\exp\left\{i\frac{a^{2}t}{8\hbar c^{3}}\hat{p}\hat{x}^{2}\right\} \\
&\quad \times \exp\left\{\frac{a^{2}t}{8c^{3}}\hat{x}\right\}\exp\left\{-i\frac{3\hbar a^{2}t}{32c^{3}}\hat{p}^{-1}\right\}
\end{aligned} \tag{9}$$

where the exponentials are correct to order a^{2}. The proof of this relation is analogous to that of the Baker-Campbell-Hausdorf theorem. For instance, an expression like $F(t) := \exp[it(\lambda\hat{p} + \lambda'\hat{p}\hat{x}^{2})]$ can be decomposed by making the ansatz $F(t) = \exp[iu(t)\hat{p}]\exp[iv(t)\hat{p}\hat{x}]\exp[iw(t)\hat{p}\hat{x}^{2}]$ and differentiating both expressions for $F(t)$ with respect to t. Using simple commutation relations the resulting equation corresponds to a set of ordinary differential equations for $u(t)$, $v(t)$, and $w(t)$ which can be solved exactly or, as is sufficient for our purposes, to order a^{2}.

Eq. (9) allows us to study in detail the time evolution of a quasi-particle mode $\Theta(x,t)$ by discussing the action of $\exp[\pm i\hat{\omega}_{+}t]$ on $|\Theta_{0}\rangle$ [7], with $\langle x|\Theta_{0}\rangle := \Theta_{0}(x)$. Before doing so we remark that for $a = 0$ we have $\hat{\omega}_{-} = \hat{\omega}_{+}$ and $\exp[\pm i\hat{\omega}_{+}t] = \exp[\pm i\hat{p}ct/\hbar]$ (in accordance with the discussion given above $p > 0$ has been assumed). The r.h.s. of this expression corresponds to a shift of the position, $x' = x \pm ct$. Hence, for $a = 0$ the exponentials $\exp[\pm i\hat{\omega}_{+}t]$ describe the propagation of sound waves to the right or to the left, respectively. As the right-moving part can be obtained from the left-moving part by time-inversion we will discuss the -left moving part $\Theta_{L}(x,t) := \langle x|\exp[-i\hat{\omega}_{+}t]|\Theta_{0}\rangle$ only.



FIG. 1. Sketch of the evolution of a localized quasi-particle mode in the linear regime of the combined potential $V(x)$. A small excitation Θ_{0} of the ground state ψ_{g} is splitted into two counterpropagating sound waves. The homogeneous force induces a squeezing operation on these two wavepackets. The additional potential provided by ψ_{g} leads to a reduction of the acceleration induced by the force.

To get an explicit expression for $\Theta_{L}(x,t)$ we have to analyze the action of each individual exponential of Eq. (9). The first one is just a c-number and corresponds to a rescaling of the wavefunction (note that Eq. (3) does not conserve the norm of Θ). As discussed above the second exponential corresponds to a time-dependent shift of the wavefunction and describes an accelerated sound propagation:

$$\langle x|\exp\left\{-i\frac{ct}{\hbar}\left(1+\frac{at}{4c}\right)\hat{p}\right\}|\phi\rangle = \langle x - ct - \tfrac{1}{4}at^2|\phi\rangle \ . \tag{10}$$

In this equation $|\phi\rangle$ denotes an arbitrary element of the Hilbert-space $\mathrm{L}^2(\mathbf{R})$ of square integrable functions. The third exponential is of the form $\exp[-i\alpha\hat{p}\hat{x}]$ with $\alpha := at[1-at/(4c)]/(2\hbar c)$. This operator corresponds to a squeezing transformation,

$$\langle x|e^{-i\alpha\hat{p}\hat{x}}|\phi\rangle = e^{-\hbar\alpha}\langle xe^{-\hbar\alpha}|\phi\rangle \ . \tag{11}$$

A quick way to see this is to write the position and momentum operator in terms of creation and annihilation operators as for the harmonic oscillator: $\hat{x}\sim\hat{a}+\hat{a}^\dagger$, $\hat{p}\sim i(\hat{a}^\dagger-\hat{a})$. The exponential then can be written as $\exp\{\alpha[(\hat{a}^\dagger)^2-\hat{a}^2-1]\}$ which is just the usual form of a single-mode squeezing transformation (see, e.g., Ref. [8]). Mathematically, Eq. (11) can be proven by differentiating $f(\alpha,x) := \langle x|e^{-i\alpha\hat{p}\hat{x}}|\phi\rangle$ with respect to α. the resulting differential equation, $\partial f/\partial\alpha = -\hbar\partial(xf)/\partial x$ has, for the inital value $f(0,x) = \langle x|\phi\rangle$, the solution (11). Analogously one can show that in momentum space the squeezing operator acts like

$$\langle p|e^{-i\alpha\hat{p}\hat{x}}|\phi\rangle = \langle pe^{\hbar\alpha}|\phi\rangle \ . \tag{12}$$

As usual the exponent of the squeezing factor $\exp[\hbar\alpha]$ has the opposite sign compared to the squeezing of the conjugate variable $\hat{x}$. New compared to the usual squeezing of ordinary particles is the c-number factor of $\exp[-\hbar\alpha]$ appearing on the r.h.s. of Eq. (11). This factor and the resulting asymmetry between Eqs. (11) and (12) is a consequence of the operator ordering: $\hat{p}\hat{x}$ instead of $(\hat{p}\hat{x}+\hat{x}\hat{p})/2$. The order of $\hat{p}\hat{x}$ itself is a consequence of the asymmetric definition of $\hat{\omega}_+ = [\hat{p}^2(c^2+a\hat{x})]^{1/2}$.

In an analogous treatment one can show that the fourth exponential, which is of the form $\exp[i\beta\hat{p}\hat{x}^2]$ with $\beta := a^2t/(8\hbar c^3)$, corresponds to a rational transformation of x,

$$\langle x|e^{i\beta\hat{p}\hat{x}^2}|\phi\rangle = \frac{1}{(1-\hbar\beta)^2}\left\langle\frac{x}{1-\hbar\beta x}\right|\phi\right\rangle \ . \tag{13}$$

The singularity at $x = 1/(\hbar\beta)$ is unphysical since it only appears if the wavefunction is not localized inside the area where $c^2 \gg ax$ holds. The corresponding equation in momentum space is much more complicated and difficult to interpret. We can gain some insight into this case by considering the Heisenberg-picture evolution of the momentum operator:

$$e^{-i\beta\hat{p}\hat{x}^2}\hat{p}e^{i\beta\hat{p}\hat{x}^2} = \hat{p}(1+\hbar\beta\hat{x})^2 \ . \tag{14}$$

Since the momentum operator gets multiplied with the conjugate operator $\hat{x}$ its action in momentum space is difficult to interpret. However, it still remains an operator linear in $\hat{p}$ so that its action in position space is easy to understand: the wavefunction gets multiplied with $(1+\hbar\beta x)^2$ and is then differentiated.

The fifth exponential in Eq. (9) acts as a position-dependent rescaling of the wavefunction. In momentum space it corresponds to a *imaginary* shift of the momentum. The last exponential, which is of the form $\exp[-i\gamma\hat{p}^{-1}]$ with $\gamma := 3\hbar a^2t/(32c^3)$, does not change the momentum of the wavefunction but affects the quasi-particle's position according to $\exp[-i\gamma\hat{p}^{-1}]\hat{x}\exp[i\gamma\hat{p}^{-1}] = \hat{x}+\gamma\hbar\hat{p}^{-2}$. This expression is not very insightful, however, so that we exploit our assumption that the quasi-particle mode is initially localized around some momentum p_0. We then can expand the operator $\hat{p}^{-1}$ as $1/p_0-(\hat{p}-p_0)/p_0^2+(\hat{p}-p_0)^2/p_0^3+\cdots$ so that we essentially get expressions which are linear or quadratic in $\hat{p}$. The linear part shifts the position analogous to Eq. (10). The term quadratic in $\hat{p}$ has the same structure as the operator $\hat{p}^2/(2M)$ of the kinetic energy of an ordinary Schrödinger particle. Hence an acceleration introduces a momentum-dependent mass $\propto p_0^3/a^2$ for sound waves in a BEC.

Putting everything together by applying successively all exponentials of Eq. (9) to the quasi-particle mode we arrive at

$$\langle x|e^{-i\hat{\omega}_+t}|\Theta_0\rangle = \left\{1-\frac{3at}{4c}+\frac{3a^2tx}{8c^3}+\frac{3a^2t^2}{32c^2}\right\} \times\exp\left[-i\frac{3a^2t\hbar}{16c^3p_0}\right]\langle x_H(t)|\Theta_0\rangle \tag{15}$$

with

$$x_H(-t) := \left(1-\frac{at}{2c}\right)x+\frac{1}{4}at^2-ct\left(1-\frac{3a^2\hbar^2}{32c^4p_0^2}\right) +\frac{a^2t}{8c^3}x^2+O(a^3)+O(1/p_0^3) \ . \tag{16}$$

This expression coincides with the Heisenberg-picture position operator $\hat{x}_H(-t) = \exp[-i\hat{\omega}_+t]\hat{x}\exp[i\hat{\omega}_+t]$ if the variable x on the r.h.s. of Eq. (16) is replaced by the position operator $\hat{x}$.

In view of the discussions given above the result (15) is easy to understand. The prefactor in curly brackets corresponds to a position dependent rescaling of the wavefunction, a combined effect of the first exponential in Eq. (9) and the prefactors present in Eqs. (11) and (13). The phase factor stems from the last exponential. In

$x_H(-t)$ the first term is a result of the squeezing transformation (11). The second term describes an acceleration of the quasi-particle motion. Since the quasi-particle has to overcome the potential created by the accelerated ground-state, which according to Eq. (2) has the opposite sign as V itself, the acceleration is only half as large as for a usual particle (prefactor 1/4 instead of 1/2). The term proportional to ct corresponds to the sound propagation, where a correction $\propto a^2/p_0^2$ to the velocity of sound appears because of the last exponential in Eq. (9). The last term, which is nonlinear in x, is caused by the rational transformation of Eq. (13).

It is also interesting to study the momentum operator $p_H(-t) := \exp[-i\hat{\omega}_+ t]\hat{p}\exp[i\hat{\omega}_+ t]$ in the Heisenberg picture, which is given by

$$p_H(-t) = \hat{p}\left(1 + \frac{at}{2c} + \frac{a^2t^2}{4c^2}\right) - \frac{a^2t}{4c^3}\hat{p}\hat{x} + i\hbar\frac{a^2t}{8c^3} + O(a^3) \tag{17}$$

The first term describes, for $a < 0$, a squeezing of the momentum. The second term has its origin in the rational transformation (14) which also produces a higher order correction to the first term. The last term is produced by the fifth exponential in Eq. (9). Because of this term $p_H(-t)$ is not a hermitean operator anymore. This is also a consequence of the asymmetric occurence of $\hat{x}$ and $\hat{p}$ in the definition of $\hat{\omega}_+$.

For an ordinary accelerated particle the momentum in the Heisenberg picture is given by $\hat{p} + Mat$ where the last term corresponds to the increase of the momentum due to the acceleration. Notably this term is absent in the case (17) of a quasi-particle. Thus, the momentum of a quasi-particle is squeezed rather than accelerated by a weak external homogeneous force.

This behaviour can be understood if we consider $-\hat{\omega}_+$ of Eq. (7) as a kind of Hamiltonian for a classical system, i.e., $H = -p[c^2 + ax]^{1/2}$ (the minus sign arises because we are dealing with $x_H(-t)$ instead of $x_H(t)$). It then becomes obvious that the classical canonical momentum is not proportional to dx/dt anymore so that its time evolution is more complicated. This is reflected by the squeezing transformation. We also remark that the results (16) and (17) also serve as a further justification for Eq. (8) since for $\hbar = 0$ they do agree with the solutions of the classical equations of motion, $\dot{x} = \partial H/\partial p$ and $\dot{p} = -\partial H/\partial x$.

Up to now we have analysed the time evolution of the left-moving wavepacket $\Theta_L(x,t) := \langle x|\exp[-i\hat{\omega}_+ t]|\Theta_0\rangle$. The corresponding right-moving part $\Theta_R(x,t) := \langle x|\exp[i\hat{\omega}_+ t]|\Theta_0\rangle$ can be obtained by reversing the sign of t in the results (15), (16), and (17). It then becomes obvious that (for $a > 0$) the momentum of Θ_R is squeezed instead of its position (see Fig. 1).

In conlusion we have demonstrated that a weak homogeneous force acting on a BEC alters the motion of quasi-particle modes in a different way than for ordinary particles. The acceleration of the position is smaller since the mode has to overcome the additional potential provided by the ground state. The momentum is squeezed rather than accelerated because the “Hamiltonian” $\hat{\omega}_+$ contains a term $\propto \hat{p}\hat{x}$ which acts in the same way as a squeezing operator in quantum optics.

Acknowledgement: The work has been supported by the Australian Research Council.

---

[1] M. Anderson, *et al.*, Science **269**, 198 (1995); C. C. Bradley *et al.*, Phys. Rev. Lett.**75**, 1687 (1995); M.-O. Mewes *et al.*, Phys. Rev. Lett. **77**, 416 (1996);

[2] D.S. Jin *et al.*, Phys. Rev. Lett. **77**, 420 (1996).

[3] D.M. Stamper-Kurn *et al.*, MIT, preprint cond-mat/9801262.

[4] V.M. Pérez-García *et a;.*, Phys. Rev. Lett. **77**, 5320 (1996); Y. Castin and R. Dum, Phys. Rev. Lett. **79**, 3553 (1997); M. Edwards *et al.*, Phys. Rev. Lett. **77**, 1671 (1996).

[5] G. Baym and C.J. Pethick, Phys. Rev. Lett. **76**, 6 (1996); E. Lundh, C.J. Pethick, and H. Smith, Phys. Rev. A **55**, 2126 (1997).

[6] A.L. Fetter, Ann. Phys. (NY) **70**, 67 (1972).

[7] The introduction of the ket $|\Theta_0\rangle$ is just a mathematical trick so that we can analyze the action of $\exp[\pm i\hat{\omega}_+ t]$ in terms of an operator acting on some Hilbert-space. $|\Theta_0\rangle$ is *not* an element of a Hilbert space for quasi-particle modes. Rather, $\Theta_0(x)$ is interpreted as a wavefunction in the Hilbert-space $\mathrm{L}^2(\mathbf{R})$ of square-integrable functions so that we can use the usual operator algebra in the form $\langle x|\hat{O}(\hat{x},\hat{p})|\Theta_0\rangle = \hat{O}(x, -i\hbar\partial/\partial x)\langle x|\Theta_0\rangle$.

[8] D.F. Walls and G.J. Milburn, *Quantum Optics*, Springer-Verlag, Berlin 1994.